arXiv:1304.0285v1 [math.CO] 1 Apr 2013

Strong chromatic index of k-degenerate graphs

Tao Wang [a, b, *]
[a]Institute of Applied Mathematics
Henan University, Kaifeng, 475004, P. R. China
[b]College of Mathematics and Information Science
Henan University, Kaifeng, 475004, P. R. China

April 2, 2013

Abstract

An edge coloring is called a *strong edge coloring* if any path with three edges has three distinct colors. The *strong chromatic index* $\chi'_s(G)$ of a graph G is the least number of colors needed in any acyclic edge coloring of G. In this paper, we show that the strong chromatic index of a k-degenerate graph G is at most $(4k-2)\cdot\Delta(G)-2k^2+1$. If G is a graph such that all of its 3^+-vertices induce a subgraph without cycles, then $\chi'_s(G) \leq 4\Delta(G) - 3$; as a corollary, every minimally 2-connected graph G has strong chromatic index at most $4\Delta(G) - 3$.

1 Introduction

An edge coloring is called a *strong edge coloring* if any path with three edges has three distinct colors. In other words, an edge coloring is a strong edge coloring if and only if for any edge wv, the color of wv is distinct with the colors of the edges (except wv) incident with $N_G(w) \cup N_G(v)$. The *strong chromatic index* $\chi'_s(G)$ of a graph G is the least number of colors needed in any acyclic edge coloring of G. The *degree* of a vertex v in G, denoted by $\deg(v)$, is the number of incident edges of G. A vertex of degree k is called a k-vertex, and a vertex of degree at most or at least k is call a k^-- or k^+-vertex, respectively.

In 1985, Edrös and Nešetřil construct graphs with strong chromatic index $\frac{5}{4}\Delta^2$ when Δ is even, $\frac{1}{4}(5\Delta^2-2\Delta+1)$ when Δ is odd. Inspired by this construction, they proposed the following strong edge coloring conjecture:

Conjecture 1. If G is a graph with maximum degree Δ, then

$$\chi'_s(G) \leq \begin{cases} \frac{5}{4}\Delta^2, & \text{if } \Delta \text{ is even} \\ \frac{1}{4}(5\Delta^2 - 2\Delta + 1), & \text{if } \Delta \text{ is odd} \end{cases}$$

Faudree et al. posed another various conjectures on the cubic graphs, for the details see [4]. A graph is *k-degenerate* if every subgraph has a vertex of degree at most k. Chang and Narayanan [2] showed the strong chromatic index of a 2-degenerate graph G is at most $10\Delta(G) - 10$. Recently, Luo and Yu [5] improved the upper bound to $8\Delta(G)-4$. For general k-degenerate graphs, Dębski et al. [3] present an upper bound $(4k-1)\Delta(G)-k(2k+1)+1$. In this note, we use the method developed in [3] and improve the upper bound to $(4k-2)\cdot\Delta(G)-2k^2+1$.

2 Result

A vertex v in a k-degenerate graph is *nice* if it is adjacent to at most k vertices of degree large than k.

Lemma 1 ([2]). If G is a k-degenerate graph, then there exists a vertex w such that at least $\max\{1, \deg(w) - k\}$ of its neighbors are k^--vertices.

Theorem 2.1. If G is a k-degenerate graph with $k \leq \Delta(G)$, then $\chi'_s(G) \leq (4k-2)\cdot\Delta(G) - 2k^2 + 1$.

Proof. By the definition of k-degenerate graph, every subgraph of G is also a k-degenerate graph. We want to obtain a sequence $\Lambda_1, \dots, \Lambda_m$ of subsets of edges as follows. For convenience, let $\Lambda_0 = \emptyset$. Suppose that Λ_{i-1} has been well defined, let G_i be the graph induced by $E(G) - (\Lambda_0 \cup \cdots \cup \Lambda_{i-1})$. Notice that G_i is empty or G_i is a

---

[*] Corresponding author: `wangtao@henu.edu.cn`

k-degenerate graph. For convenience, denote the degree of v in G_i by $\deg_i(v)$. If G_i is nonempty, then choose a nice vertex w_i of G_i as described in Lemma 1, and let

$$\Lambda_i = \{w_i v_i \mid \deg_i(v_i) \leq k\}.$$

By Lemma 1, it follows that $\Lambda_i \neq \emptyset$ and this process will terminate with a subset L_m. Notice that the subgraph induced by Λ_i is a star in the graph G.

Claim 1. There exists no $j < i$ such that $w_i = w_j$.

Proof. Suppose to the contrary that there exists $j < i$ such that $w_i = w_j$. From the construction, we have that $\deg_{j+1}(w_i) \leq k$ and $\deg_j(v_i) > k$. But $\deg_i(v_i) \leq k$, thus there exists $j < t < i$ such that v_i is a nice vertex in G_t but $\deg_t(w_i) > k$, which leads to a contradiction that $\deg_{j+1}(w_i) \leq k < \deg_t(w_i)$. □

We will color the edges from Λ_m to Λ_1, we remind the readers this is the reverse order of the ordinary sequence.

In the following, we want to give an algorithm to obtain a strong edge coloring of G. First, we can color the edges in Λ_m with distinct colors. Now, it is the edge $w_i v_i$'s turn to color, where $w_i v_i \in \Lambda_i$. In order to guarantee the result coloring is a partial strong edge coloring in any step, the two edges which are incident with $N_G(w_i) \cup N_G(v_i)$ must receive distinct colors.

Now, we compute the number of colored edges in G_i which are incident with $N_G(w_i) \cup N_G(v_i)$. Let x_i be the number of neighbors of w_i outside of G_i, let y_i be the number of neighbors of w_i with degree at most k in G_i, and let z_i be the number of neighbors of w_i with degree greater than k in G_i. Suppose that $x_i > 0$ and let w be an arbitrary neighbor of w_i outside G_i. By the claim, the edge ww_i is in some Λ_t with center w, thus $\deg_t(w_i) \leq k$ and w is incident with at most k colored edges. Furthermore, it implies that $\deg(w_i) \leq k$. Therefore, if $x_i \neq 0$, then the number of edges which are incident with $N_G(w_i) \setminus \{v_i\}$ is at most

$$\begin{aligned}&(x_i + y_i - 1) \cdot k + z_i \cdot \Delta(G)\\ =&(x_i + y_i - 1 + z_i) \cdot k + z_i \cdot (\Delta(G) - k)\\ \leq&(k-1)k + (k-2) \cdot (\Delta(G) - k), \quad \text{note that } x_i + y_i + z_i \leq k \text{ and } z_i \leq k-2;\end{aligned}$$

if $x_i = 0$, then the number of edges which are incident with $N_G(w_i) \setminus \{v_i\}$ is at most

$$\begin{aligned}&(y_i - 1) \cdot k + z_i \cdot \Delta(G)\\ =&(y_i - 1 + z_i) \cdot k + z_i \cdot (\Delta(G) - k)\\ \leq&(\Delta(G) - 1) \cdot k + k \cdot (\Delta(G) - k), \quad \text{note that } z_i \leq k.\end{aligned}$$

Let p_i be the number of neighbors of v_i outside of G_i and q_i be the number of neighbors in G_i. If $\deg(v_i) > k$, then $p_i > 0$ and there exists some j with $j < i$ such that v_i is the center of Λ_j. It follows that $\deg_{j+1}(v_i) \leq k$ and for every edge $v_i v$ in Λ_j, the vertex v is incident with at most $k-1$ colored edges. Therefore, if $\deg(v_i) > k$, then the number of colored edges which are incident with $N_G(v_i) \setminus \{w_i\}$ is at most

$$\begin{aligned}&(k-1) \cdot (\deg(v_i) - \deg_{j+1}(v_i)) + (\deg_{j+1}(v_i) - 1) \cdot \Delta(G)\\ =&(k-1) \cdot \deg(v_i) + \deg_{j+1}(v_i) \cdot (\Delta(G) - k + 1) - \Delta(G)\\ \leq&(k-1)\Delta(G) + k \cdot (\Delta(G) - k + 1) - \Delta(G), \quad \text{note that } \deg_{j+1}(v_i) \leq k;\end{aligned}$$

if $\deg(v_i) \leq k$, then the number of colored edges which are incident with $N_G(v_i) \setminus \{w_i\}$ is at most $(k-1) \cdot \Delta(G)$.

Hence, the number of colored edges incident with w_i or v_i is at most $(4k-2) \cdot \Delta(G) - 2k^2$, and there are at least one available color for $w_i v_i$. When all the edges are colored, we obtain a strong edge coloring of G. □

Corollary 1. If G is 2-degenerate graph, then $\chi'_s(G) \leq 6\Delta(G) - 7$.

Next, we investigate a class of graphs whose all 3^+-vertices induce a subgraph without cycles.

Lemma 2. If G is a graph with at least one edge such that all the 3^+-vertices induce a subgraph without cycles, then there exists a vertex w such that at least $\max\{1, \deg_G(v) - 1\}$ of its neighbors are 2^--vertices.

Proof. Let A be the set of 2^--vertices. It is obvious that A is nonempty, for otherwise every vertex has degree at least three, and then G contains cycles. Now, consider the graph $G - A$. If $G - A$ is empty, then every nonisolated vertex satisfies the desired. So we may assume that the $G - A$ is nonempty. Since the graph $G - A$ is a forest, it follows that there exists at least one 1^--vertices in $G - A$, and every such vertex satisfies the desired. □

Theorem 2.2. If G is a graph such that all of its 3^+-vertices induce a subgraph without cycles, then $\chi_s'(G) \leq 4\Delta - 3$.

Proof. Let H be an arbitrary subgraph with at least one edge, it is obvious that all of its 3^+-vertices also induce a subgraph without cycles. We want to obtain a sequence $\Lambda_1, \ldots, \Lambda_m$ of subsets of edges as follows. For convenience, let $\Lambda_0 = \emptyset$. Suppose that Λ_{i-1} has been well defined, let G_i be the graph induced by $E(G) - (\Lambda_0 \cup \cdots \cup \Lambda_{i-1})$. Notice that G_i is empty or G_i is a graph such that all of its 3^+-vertices induce a subgraph without cycles. For convenience, denote the degree of v in G_i by $\deg_i(v)$. If G_i is nonempty, then choose a *nice vertex* w_i of G_i as described in Lemma 2, in other words, at least $\max\{1, \deg_i(w_i) - 1\}$ of w_i's neighbors are 2^--vertices; and let

$$\Lambda_i = \{w_i v_i \mid \deg_i(v_i) \leq 2\}.$$

By Lemma 2, it follows that $\Lambda_i \neq \emptyset$ and this process will terminate with a subset L_m. Notice that the subgraph induced by Λ_i is a star in the graph G.

Claim 2. There exists no $j < i$ such that $w_i = w_j$.

Proof. Suppose to the contrary that there exists $j < i$ such that $w_i = w_j$. From the construction, we have that $\deg_{j+1}(w_i) \leq 2$ and $\deg_j(v_i) > 2$. But $\deg_i(v_i) \leq 2$, thus there exists $j < t < i$ such that v_i is a nice vertex in G_t but $\deg_t(w_i) > 2$, which leads to a contradiction that $\deg_{j+1}(w_i) \leq 2 < \deg_t(w_i)$. □

We will color the edges from Λ_m to Λ_1, we remind the readers this is the reverse order of the ordinary sequence.

In the following, we want to give an algorithm to obtain a strong edge coloring of G. First, we can color the edges in Λ_m with distinct colors. Now, it is the edge $w_i v_i$'s turn to color, where $w_i v_i \in \Lambda_i$. In order to guarantee the result coloring is a partial strong edge coloring in any step, the two edges which are incident with $N_G(w_i) \cup N_G(v_i)$ must receive distinct colors.

Now, we compute the number of colored edges in G_i which are incident with $N_G(w_i) \cup N_G(v_i)$. Let x_i be the number of neighbors of w_i outside of G_i, let y_i be the number of neighbors of w_i with degree at most 2 in G_i, and let z_i be the number of neighbors of w_i with degree greater than k in G_i. Suppose that $x_i > 0$ and let w be an arbitrary neighbor of w_i outside G_i. By the claim, the edge ww_i is in some Λ_t with center w, thus $\deg_t(w_i) \leq 2$ and w is incident with at most one colored edges. Furthermore, it implies that $x_i = y_i = 1$ and $z_i = 0$. Therefore, if $x_i \neq 0$, then the number of edges which are incident with $N_G(w_i) \setminus \{v_i\}$ is at most one; if $x_i = 0$, then the number of edges which are incident with $N_G(w_i) \setminus \{v_i\}$ is at most

$$\begin{aligned}
&(y_i - 1) \cdot 2 + z_i \cdot \Delta(G)\\
=&2(y_i - 1 + z_i) + z_i \cdot (\Delta(G) - 2)\\
\leq&3\Delta(G) - 4, \quad \text{note that } z_i \leq 1;
\end{aligned}$$

Let p_i be the number of neighbors of v_i outside of G_i and q_i be the number of neighbors in G_i. If $\deg(v_i) > k$, then $p_i > 0$ and there exists some j with $j < i$ such that v_i is the center of Λ_j. It follows that $\deg_{j+1}(v_i) = 1$ and for every edge $v_i v$ in Λ_j, the vertex v is incident with at most one colored edges. Therefore, if $\deg(v_i) > 2$, then the number of colored edges which are incident with $N_G(v_i) \setminus \{w_i\}$ is at most $\deg_G(v_i) - 1 \leq \Delta(G) - 1$; if $\deg(v_i) \leq 2$, then the number of colored edges which are incident with $N_G(v_i) \setminus \{w_i\}$ is at most $\Delta(G)$.

Hence, the number of colored edges incident with w_i or v_i is at most $4\Delta(G) - 4$, and there are at least one available color for $w_i v_i$. When all the edges are colored, we obtain a strong edge coloring of G. □

A 2-connected graph G is *minimally* 2-*connected* if $G - e$ is not 2-connected for any edge e in G.

Theorem 2.3 ([1]). The minimum degree of a minimally 2-connected graph is two, and the subgraph induced by all the 3^+-vertices has no cycles.

Corollary 2. If G is a minimally 2-connected graph, then $\chi_s'(G) \leq 4\Delta(G) - 3$.

Remark 1. This bound is sharp since $\chi_s'(C_5) = 5 = 4\Delta(C_5) - 3$.

A graph is *chordless* if every cycle is an induced cycle. It is easy to show that a 2-connected graph is chordless if and only if it is minimally 2-connected. Every chordless graph is a 2-degenerate graph. In [2], Chang and Narayanan proved the strong chromatic index of a chordless graph is at most $8\Delta(G) - 6$, and Dębski et al. [3] improved the upper bound to $4\Delta(G) - 3$, but I doubt that the proofs are correct since they used an incorrect lemma (see [2, Lemma 7]).

References

[1] B. Bollobás, Extremal graph theory, Dover Publications Inc., Mineola, NY, 2004, reprint of the 1978 original.

[2] G. J. Chang and N. Narayanan, Strong chromatic index of 2-degenerate graphs, J. Graph Theory (2013).

[3] M. Dębski, J. Grytczuk and M. Śleszyńska-Nowak, Strong chromatic index of sparse graphs, arXiv:1301.1992 (01/2013).

[4] R. J. Faudree, R. H. Schelp, A. Gyárfás and Z. Tuza, The strong chromatic index of graphs, Ars Combin. 29 (1990) (B) 205–211.

[5] R. Luo and G. Yu, A note on strong edge-colorings of 2-degenerate graphs, arXiv:1212.6092 (12/2012).